

#1 ALL PURPOSE
HEAVY DUTY
CLEANER

Spray X Disinfectant is a broad-spectrum virucidal hard surface disinfectant that is expected to inactivate the SARS-CoV-2 (The virus that causes COVID-19).



Cleaner • Deodorizer • Disinfectant • Kills Mold
Nettoyant • Désodorisant • Désinfectant • Antimoisissure



DIN# 02500426

Based on laboratory testing Spray X Disinfectant demonstrated effectiveness against SARS Associated Coronavirus in 2 minutes on commonly touched surfaces such as a car interior, door handles, tools and equipment. For details on different viruses contact time see page 4.

1 (844) 500 2436
www.rexalllabs.com
445 Eastchester Ave E St.
Catharines, ON Canada L2M 6S2

Rexall
LABORATORIES & CHEMICALS

Spray X



A ready-to-use, quaternary ammonium disinfectant spray cleaner that is ideal for healthcare facilities, cannabis facilities, food processing facilities, schools due to its extensive list of disinfectant claims including tuberculosis, SARS associated coronavirus and norovirus. Inhibits growth of mold and mildew and controls related odors. Health Canada registered as a broad spectrum disinfectant cleaning product.

RECOMMENDED FOR:

- Hard, non-porous surfaces
- Floors
- Walls
- Counters
- Sinks
- Appliances
- Metal Surfaces
- Stainless Steel Surfaces
- Glazed Porcelain
- Ceramic Tile
- Plastic Surfaces
- Bathrooms
- Shower Stalls and Bathtubs
- Poultry Equipment
- Animal Quarters and Kennels
- Hospitals, Health Care facilities

SPECIFICATIONS:

Registrations.....**Health Canada**
Dilution.....RTU
pH Level.....11.8 – 12.8
Color.....Clear
Fragrance.....Free



DIRECTIONS:

- Remove gross filth or heavy soil prior to application of the product.
- Hold container 15-20 cm from surface to be treated. Spray area until it is covered with the solution.
- Allow product to penetrate and remain wet for 10 minutes.
- Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.
- A rinse with potable water is required for surfaces in direct contact with food. In addition, this product.

BACTERICIDAL ACTIVITY

When used as directed, this product exhibits effective disinfectant activity against the organisms: Staphylococcus aureus, Salmonella enterica, Pseudomonas aeruginosa, Escherichia coli, Escherichia coli 0157:H7, Corynebacterium ammoniagenes, Enterococcus faecium, Listeria monocytogenes, Yersenia enterocolitica, Salmonella (typhi) enterica, Vancomycin Resistant Enterococcus faecalis, Vancomycin Resistant Staphylococcus aureus and Methicillin Resistant Staphylococcus aureus, Staphylococcus aureus (VISA), Streptococcus pyogenes (Necrotizing Fasciitis), Methicillin Resistant Staphylococcus epidermidis (MRSE), Community Associated Methicillin Resistant Staphylococcus aureus (CA-MRSA), Klebsiella pneumonia New Delhi Metallo-Beta Lactamase (NDM-1) Carbapenem Resistant and meets the requirements for hospital use.

ONE-STEP CLEANING AND DISINFECTION OF TOILET BOWLS AND URINALS

Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 30-60 mL of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 10 minutes and flush.

DISINFECTION, DEODORIZING AND CLEANING

Remove gross filth or heavy soil prior to application of the product. Hold container 15-20 cm from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and remain wet for 10 minutes. No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

BROAD-SPECTRUM VIRUCIDAL ACTIVITY

This product, when used on environmental, inanimate, non-porous surfaces exhibits effective virucidal activity against HIV-1, Hepatitis C Virus (HCV), Bovine Viral Diarrhea Virus (BVDV), Human Coronavirus, Hepatitis A Virus, Norovirus (including Norwalk Virus), Feline Calicivirus, Rabies Virus, Canine Parvovirus, Poliovirus Type 1, Avian Influenza A, Turkey/Wisc/(H9N2), Avian Influenza (H3N2), Influenza A (H1N1), Paramyxovirus, Rotavirus, Rhinovirus type 39 and SARS Associated Coronavirus. This product has demonstrated effectiveness against Bovine Viral Diarrhea (BVDV), and Influenza A virus and is expected to inactivate all Influenza A viruses including 2009 (H1N1) pandemic Influenza A virus.

DISINFECTION OF POULTRY EQUIPMENT

For disinfection of pre-cleaned poultry equipment (brooders, watering founts, feeding equipment). Remove all poultry and feeds from premises, trucks, coops and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or transversed by poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for 10 minutes. Ventilate building, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried. All treated equipment must be rinsed with potable water before reuse.

DISINFECTION OF ANIMAL QUARTERS AND KENNELS

WHEN TESTED AGAINST THE PROTOCOL VIRUCIDAL EFFICACY OF DISINFECTANTS FOR USE ON INANIMATE ENVIRONMENTAL SURFACES, EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS EFFECTIVE AGAINST CANINE PARVOVIRUS IN THE PRESENCE OF ORGANIC SOIL (5% FETAL BOVINE SERUM).

For disinfection of pre-cleaned animal quarters and kennels. Remove all animals and feeds from premises, trucks and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or transversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for 10 minutes. Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment must be rinsed with potable water before reuse.

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or bodily fluids, and in which the surfaces/objects likely to be soiled with blood or bodily fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1.

Special Instructions for Cleaning and Decontamination against HIV-1 of Surfaces Soiled with Blood or Body Fluids

Personal Protection: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

Cleaning Procedures: Target surfaces must be cleaned prior to disinfection, as the presence of heavy soil on a surface may reduce the intended disinfectant efficacy of the product.

Contact Time/Concentration: Apply product directly and allow surface to remain wet for 10 minutes.

Infectious materials disposal: Any materials used in the cleaning process that may contain blood or body fluids are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

MILDEWSTAT

To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

FUNGICIDAL ACTIVITY

This product is fungicidal against the pathogenic fungi, Trichophyton mentagrophytes (Athlete's Foot Fungus) when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).



EMERGING PATHOGEN COMPLIANCE GUIDANCE

Spray X product has demonstrated effectiveness against SARS Associated Coronavirus on hard, non-porous surfaces and is on HEALTH CANADA'S list of disinfectants with evidence for use against COVID-19.

1/2 MINUTE CONTACT TIME:

- Norovirus
- Feline Calicivirus
- Rabies Virus

1 MINUTE CONTACT TIME:

- Hiv-1 [Aids Virus]

2 MINUTE CONTACT TIME:

- Avian Influenza A (H3n2 & H9n2)
- Pandemic 2009 H1n1 Influenza A Virus
- Human Coronavirus
- Sars Associated Coronavirus

3 MINUTE CONTACT TIME:

- Rotavirus
- Paramyxovirus
- Rhinovirus Type 39
- Pseudomonas Aeruginosa
- Staphylococcus Aureus
- Salmonella Enterica
- Methicillin Resistant Staphylococcus Epidermis
- Ca - Methicillin Resistant Staphylococcus Epidermis
- Bovine Viral Diarrhea Virus
- Hepatitis B

- Hepatitis C
- Corynebacterium Ammonia Genes
- Vanomycin Resistant Enterococcus Faecalis
- Escherichia Coli
- Listeria Monocytogenes
- Yersinia Enterocolitica
- Streptococcus Pyogenes
- Vancomycin Intermediate Resistant

10 MINUTE CONTACT TIME:

- Poliovirus Type 1
- Canine Parvovirus
- Trichopyton Mentagrophytes
- Hepatitis A Virus

**This is not a complete list of organisms and names.*